

NTRR - No.32

Natural Therapies Research Report

Nutra Therapeutics

Essential Fatty Acid Supplementation: IMPROVES SKIN TEXTURE PLUS ENHANCES OVERALL HEALTH STATUS

By James Meschino, D.C., M.S. Research and Clinical Director, RenaiSanté Institute of Integrative Medicine

INTRODUCTION

An increasing number of holistic practitioners recommend essential fatty acid supplementation to their patients as means to enhance their overall health status or as part of the complementary treatment for various health conditions. For example, many health practitioners recommend essential fatty acid supplementation for the prevention of heart disease and cancer. With respect to the prevention of heart disease, studies suggest that omega-3 fats (from fish and flaxseed oil) serve as precursors to the formation of the type of prostaglandin hormones (PG-3) that dilate blood vessels, reduce inflammation of the arterial wall and inhibit abnormal platelet coagulation, all of which are associated with a reduced incidence of cardiovascular events. Cancer studies also show that the prostaglandin hormones made from omega-3 fats reduce the rate of cellular proliferation, which in turn, reduces the chances of genetic mutations and the development of some cancers. As well, supplementation with essential oils that contain gamma-linolenic acid (GLA) and/or omega-3 fats have been shown to be useful in the management of arthritis and inflammatory conditions, diabetic neuropathy, eczema, psoriasis, PMS, fibrocystic breast disease, menopausal symptoms, attention deficit disorder, and other conditions.

PROSTAGLANDIN SERIES 2 (PG-2) THE UNDESIRABLE PROSTAGLANDIN

Within epidermal cells PG-2 is formed from the unsaturated fat known as arachidonic acid, which is found in rich concentrations in high fat meats and high fat dairy products. As well, the overconsumption of linoleic acid (an unsaturated fat found in corn oil. sunflower seed oil and safflower seed oil, and mixed vegetable oils) encourages the conversion of linoleic acid to arachidonic acid by the body, adding to the cell membrane concentrations of arachidonic acid. Higher cell membrane concentrations of arachidonic acid, from the over-consumption of these foods, tends to favor the synthesis of PG-2, contributing to poor skin texture and appearance. Unfortunately, the typical North American diet is a rich source of arachidonic acid and linoleic acid, and thus, most individuals produce too much PG-2 in their epidermal cells, which results in skin texture and appearance that is much less smooth and soft than is attainable, and contributes to a host of poor complexion problems and aggravates skin problems such as eczema and psoriasis. (2,4) Experimental studies also show that skin cells that exhibit a high concentration of arachidonic acid are more prone to undergoing cancerous and inflammatory changes upon exposure to ultra-violet light than are skin cells that have higher concentrations of omega-3 fats. This appears to be due to the fact that the conversion of arachidonic acid to PG-2 produces an effect whereby skin cells divide at a much faster rate, which makes them more prone to cancerous mutations and inflammatory responses. Conversely, omega-3 fats, from flaxseed oil and fish oil, have been shown to slow the division rate of skin cells via their effects on promoting the formation of PG-3 which, in turn, reduces the of mutations and inflammatory incidence response by skin cells even when exposed to ultra-violet light. Thus, omega-3 fat supplementation is not only an important intervention to improve skin texture, but may help guard against the development of skin cancer and slow skin aging. (5)

PROSTAGLANDIN SERIES 1 (PG-1)

In contrast to the deleterious effects of PG-2 on the skin, prostaglandin series 1 (PG-1) and prostaglandin series 3 (PG-3) make the skin soft, smooth, silky and moist. The key building block for PG-1 is an unsaturated fat known as GLA (gamma-linolenic acid), which is found in high concentrations in borage oil (22% yield, whereas evening primrose oil is only a 9% yield). GLA can also be formed in the body from linoleic acid, but individuals with certain conditions (diabetes, eczema) have been shown to have a defect in the enzyme that converts linoleic acid to GLA (delta-6 desaturase enzyme). As well, the consumption of alcohol, refined sugars, and hydrogenated fats tend to inhibit the conversion of linoleic acid to GLA. The aging process itself also slows the conversion of linoleic acid to GLA as the delta-6 desaturase enzyme becomes more sluggish. As such, most, if not all, individuals have sub-optimal cell membrane concentrations of GLA, and thus, GLA supplementation (e.g. borage oil) has been shown to improve skin texture and various skin conditions (eczema) in human studies via its conversion to PG-1 within epidermal cells. (2,6,7,8,9) Also note that the commonly seen sub-optimal intake of Vitamin B6, zinc and magnesium slows the conversion of linoleic acid to GLA, as these nutrients act as coenzymes in this biochemical reaction. (2,3)

PROSTAGLANDIN SERIES 3 (PG-3)

PG-3 also makes the skin very smooth, soft, silky and moist. PG-3 is formed from the omega-3 unsaturated fat known as EPA (eicosapentaenoic acid), which is found in cold-water marine fish such as salmon, mackerel, anchovies, sardines, and tuna, and supplements containing EPA. The body can also convert the omega-3 unsaturated fat ALA (alpha-linolenic acid) into EPA, increasing the production of PG-3. ALA is found in rich concentrations in flaxseed oil (58% yield). Thus, supplementation with flaxseed oil and/or a highyield fish oil have been shown to significantly increase production of PG-3, improving skin texture and appearance, as well as certain skin conditions (psoriasis and eczema). Fish oil also contains DHA (docosahexaenoic acid), which the body can convert into EPA and thus, PG-3, if required. DHA is also used to promote the development and function of the brain and is required for vision. (2,3,10,11,12)

PG-3 is considered to be very important for total body wellness as it also reduces risk of heart attacks by dilating blood vessels and reducing abnormal blood clotting. PG-3 has been shown to reduce cancer risk by slowing down the cell division rates (more rapid cell division leads to increased genetic mutations and thus, increased formation of cancer cells, with less time for DNA repair enzymes to correct the mistakes). (2) This has also been shown to be true with respect to skin cancer in experimental studies, whereby higher PG-2 levels in skin cells produced a significantly higher yield of cancer development upon exposure to ultra-violet light. PG-3 is also known to reduce inflammation (including skin inflammatory responses), a role it shares with PG-1. (5) Thus, supplementation with borage oil, flaxseed oil and a high-yield fish oil (30% EPA/20% DHA) is not only beneficial to improve skin texture and appearance, and to treat certain skin conditions, but it is also of value in the global prevention of heart and cardiovascular disease, cancer, and in the management of inflammatory conditions such as arthritis, Crohn's disease, colitis, diabetic neuropathies, skin inflammatory conditions, etc. (2)

VITAMINS AND MINERALS AS CO-FACTORS FOR PROSTAGLANDIN SYNTHESIS

It should be noted that certain vitamins and minerals are required as co-factors in the enzymatic reactions that allow skin cells to convert essential oils (flaxseed, borage and fish oils) into PG-1 and PG-3. For example, the conversion of ALA to EPA requires optimal nutritional status of Vitamin B6, zinc, magnesium, and niacin (Vitamin B3), as coenzymes. As well, the synthesis of PG-1 and PG-3 also requires optimal intake of Vitamin C, Vitamin E, and selenium. These antioxidants affect the enzyme cyclooxygenase, which is the final enzyme in the conversion of essential fats to PG-1 and PG-3. Also note that supplementation with omega-3 fats (ALA, EPA and DHA) also inhibits the conversion of GLA to arachidonic acid by inhibiting the delta-5 desaturase enzyme. Thus, ALA, EPA and DHA not only increase the synthesis of PG-3, but they help to inhibit the cell membrane build up of arachidonic acid, and thus, PG-2 synthesis. (2,3)

SUMMARY AND CONCLUSION

Clinical and investigative studies now confirm that supplementation with a combination of flaxseed, borage and fish oils, at the correct dosages, is an important step to improving skin texture, complexion and the treatment of certain skin disorders (eczema, psoriasis, some cases of acne), due to their effects on promoting the prostaglandin series-1 synthesis of and prostaglandin series-3 within developing skin cells. As well, higher skin cell concentrations of omega-3 fats (from fish and flaxseed oil) may offer additional protection against ultra-violet light-induced skin cancer and photo-aging of the skin, according to emerging experimental data. Thus, essential fatty acid supplementation represents an important component of lifelong skin care management. Health care professionals should keep in mind that optimal doses of certain vitamins and minerals are required to facilitate the efficient conversion of GLA (from borage oil) to PG-1 and ALA and EPA (from flaxseed and fish oil, respectively) to PG-3. Here is a practical guide to help translate this information into a step-by-step daily formula for your patients regarding the effective use of essential fatty acid supplementation, regardless of what health or skin condition is the primary target of this intervention.

 To reduce the buildup of arachidonic acid (and thus PG-2 synthesis), avoid or restrict the intake of high fat meat and high fat dairy products (chicken, turkey and fish are good alternatives as well as non-fat milk and yogurt, and cheeses that are less than 4% milk fat). Substitute olive oil, canola oil and peanut oil in place of corn oil, sunflower seed oil, safflower seed oil and mixed vegetable oils, for salad dressings, stir-fries and to sauté vegetables. Consume alcohol in moderation, if at all, and reduce intake of refined sugars and hydrogenated fats.

- To enhance the production of PG-1 and PG-3 (which improve the softness and smoothness of the skin and help alleviate certain skin conditions), supplement the diet with an all-inone essential fatty acid supplement providing 400 mg each of flaxseed, borage and fish oils (1,200 mg capsule). For best results take 2-3 capsules per day. The fish oil should yield 30% EPA and 20% DHA content. Patients should also be encouraged to eat more fish.
- To facilitate the conversion of essential fats to PG-1 and PG-3, a high potency multi-vitamin formula should be taken that provides the following daily dosages: Vitamin C-1,000 mg; Vitamin E-400 IU (all natural); Beta-carotene-10,000 IU; Selenium-100 mcg; Zinc-15 mg; Vitamin A-2,500 IU, and a B-50 complex (as well as all other vitamins and minerals from Vitamin A to Zinc).

References

- Kaunitz, Andrew M., M.D. Use of Combination Hormone Pustisek N, Lipozencic, J. Prostaglandins in dermatology.vActa dermatovenerologica Croatica: ADC (Acta Dermatovenerol Croat) 2001 Dec; 9(4):291-8
- Murray M. The Encyclopedia of Nutritional Supplements. Prima Publishing 1996;Chapter 33:Essential Fatty Acid Supplementation:249-78
- Murray M, Pizzorno J. Encyclopedia of Natural Medicine, revised 2nd edition. Prima Publishing 1998:448-54
- Ziboh VA. Miller, C.C., Cho, Y. Metabolism of polyunsaturated fatty acids by skin epidermal enzymes: generation of antiinflammatory and antiproliferative metabolites. Am J Clin Nutr 2000 Jan;71(1 Suppl):361S-6S
- Fishcher SM. Is cyclooxygenase-2 important in skin carcinogenesis? J Environ Pathol Toxicol Oncol, 2002; 21(2):183-91
- Horrobin DF. Essential fatty acid metabolism and its modification in atopic eczema. Am J Clin Nutr, 2000 Jan;71(1 Suppl):367S-72S
- Manku MS, Horrobin DF, Morse N, Kyte V, Jenkins K, Wright S, Burton JL. Prostaglandins Leukot Med, 1982 Dec;9(6):615-28
- Raederstorff D, Loechleiter V, Moser, U. Polyunsaturated fatty acid metabolism of human skin fibroblasts during cellular aging.Int J Vitam Nutr Res 1995;65(1):51-5
- Reichert R. Evening Primrose Oil Cream, Dry Skin, & Atopic Disposition. Quarterly Review of Natural Medicine, Spring'98:p7

- Miller CC, Tang, W, Ziboh, VA, Fletcher, MP. Dietary supplementation with ehtyl ester concentrates of fish oil (n-3) and borage oil (n-6) polyunsaturated fatty acids induces epidermal generation of local putative anti-inflammatory metabolites. J Invest Dermatol 1991 Jan;86(1):96-103
- 11. Danno K, Ikai K, Imamura, S. Anti-inflammatory effects of eicosapentaenoic acid on experimental skin inflammation models. Arch Dermtol Res 1993;285(7):432-5
- Ziboh VA. Implications of dietary oils and polyunsaturated fatty acids in the management of cutaneous disorders. Arch Dermatol 1969 Feb;125(2):p241-5EPA And Skin Health